

# Wyoming Basin & Water Supply Outlook Report February 1, 2022





Photo courtesy of Wyoming Snow Survey

#### Basin Outlook Reports And Federal - State - Private Cooperative Snow Surveys

For more Wyoming water supply information, contact:

Jim Fahey - Hydrologist 100 East "B" Street, Casper, WY 82601 (307) 233-6787 james.fahey@usda.gov

#### How forecasts are made

Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snow courses and automated SNOTEL sites, along with precipitation, antecedent streamflow, and indices of the El Niño / Southern Oscillation are used in computerized statistical and simulation models to prepare runoff forecasts. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less water.

**Note**: The **median** is the official normal for snowpack (SWE), precipitation, reservoir storage, and streamflow calculations. Please refer to the **Appendix** of this report for more detailed information.


The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers. If you believe you experienced discrimination when obtaining services from USDA, participating in a USDA program, or participating in a program that receives financial assistance from USDA, you may file a complaint with USDA. Information about how to file a discrimination complaint is available from the Office of the Assistant Secretary for Civil Rights. USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex (including gender identity and expression), marital status, familial status, parental status, religion, sexual orientation, political beliefs, genetic information, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibite discrimination, I of discrimination, complete, sign, and mail a program discrimination complaint form, available at any USDA office location or online at www.ascr.usda.gov, or write to: USDA Office of the Assistant Secretary for Civil Rights 1400 Independence Avenue, SW. Washington, DC 20250-9410 Or call toll free at (866) 632-9992 (voice) to obtain additional information, the appropriate office or to request documents. Individuals who are deaf, hard of hearing, or have speech disabilities may contact USDA through the Federal Relay service at (800) 877-8339 or (800) 845-6136 (in Spanish). USDA is an equal opportunity provider, employer, and lender. Persons with disabilities who require alternative means for communication of program information (e.g., Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

### Summary

- Wyoming snowpack and/or snow water equivalents (SWEs)were below median by late January.
- Precipitation totals across Wyoming for January were **below** median. Water year precipitation totals continued to be **above** median.
- Overall reservoir storages for late January continue to be **below** median.
- Stream flow snowmelt volumes during April through July across Wyoming are forecasted to be generally near median.

#### Snowpack/SWEs

Snow water equivalents(SWEs)across Wyoming for February  $1^{st}$  were around 90% of median. SWEs in the Laramie Watershed were the highest at near 110% of median, while SWEs in the Belle Fourche and Cheyenne River Basins were the lowest at near 50% of median. Last year, SWEs across the state were 73% of median. (For complete **tabular data**, see attachment to report)




Map 1. Wyoming SWEs—Feb 1, 2022.


Wyoming Water Supply Outlook Report

#### **Precipitation**

Basin precipitation across Wyoming was near **90**% of median during January. The Laramie and Lower North Platte and South Platte River Watersheds had the <u>highest</u> precipitation totals for the month at **145** to **170**% of median. The Madison Headwaters and Cheyenne River Basins had the <u>lowest</u> precipitation amounts at near **45**% of median. Water year precipitation (October - January) is currently about **110**% of median. (For complete **tabular data**, see attachment to report)



Map 2. Current monthly precipitation by basin.



Map 3. Water year to date precipitation by basin.





Chart 1. Departure from median precipitation (water year).

#### Reservoirs

Reservoirs across Wyoming were averaging near 62% of capacity-down from 72% of capacity <u>last</u> year. Overall reservoir storages for late January are below median at 88% (101% last year). The Wind River Basin had reservoir storages at 106% of median, while the Snake River Basin had reservoir storages at 28% of median. (For complete **tabular data**, see attachment to report)



Map 4. Reservoir storage (teacup diagram) by basin.

#### **Stream Flows**

Snowmelt runoff stream flow volumes across the state are expected to be near median at around **95**%. The <u>highest</u> forecasted stream flows due to snowmelt are across the Wind, Upper North Platte, and Laramie Basins at between **110** to **120**% of median. The <u>lowest</u> snowmelt runoff volumes are expected across the Powder and Cheyenne Drainages between **65**% and **75**% of median. (For complete **tabular data**, see attachment to report)







Chart 2. 50% exceedance for major forecast points.





#### **Snake River Basin**

- The overall Snake River Basin SWE is near 95% of median.
- Last month's precipitation for the Snake River Basin was near **80**% of median. Wateryear-to-date precipitation is near **110**% of median.
- Current reservoir storage is near 28% of median for the two main reservoirs in the basin.
- The streamflow forecasts for April through July are **above** median (101%) for this basin. Jackson Lake inflows are to be 103% of median.













### Madison Headwaters Basin

- The overall basin SWE is around **95**% of median.
- Last month's precipitation for the basin was near 45% of median. Water-year-to-date precipitation is around 110% of median.






No reservoir data for the basin.


There are no streamflow forecast points for the basin.




### Yellowstone River Basin


- The overall Yellowstone River Basin SWE is near **90**% of median.
- Last month's precipitation for the Yellowstone River Basin was near **75**% of median. Water-year-to-date precipitation is near **100**% of median.
- The 50% exceedance forecasts for April through July are **below** median (**91**%) for this basin. Clarks Fork near Belfry is forecasted to have flows at **88**% of median.






No reservoir data for the basin.








### Wind River Basin

- The overall Wind River Basin SWE is near 105% of median.
- Last month's precipitation for the Wind River Basin was between **110** and **115**% of median. Water-year-to-date precipitation is around **120**% of median.
- Current reservoir storage is around 105% of median for three main reservoirs in the basin.
- The streamflow forecasts for April through July are **above** median (**115**%) for this basin. Boysen Reservoir inflows are expected to be **110**% of median.















# **Bighorn River Basin**

- The overall Bighorn River Basin SWE is near 85% of median.
- Last month's precipitation for the Bighorn River Basin was near **90**% of median. Water-year-to-date precipitation is about **105**% of median.
- Current reservoir storage is near 95% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July are <u>near</u> median (**96**%) for this basin.














### Shoshone River Basin

- The overall Shoshone River Basin SWE is around **90**% of median.
- Last month's precipitation for the Shoshone River Basin was between **85** and **90**% of median. Water-year-to-date precipitation is around **95**% of median.
- Current reservoir storage is near 85% of median for one main reservoir in the basin.
- Streamflow forecasts for April through July are **below** median (**94**%) for this basin. Buffalo Bill Reservoir inflows are expected to be **90**% of median














#### Powder River Basin

- The overall Powder River Basin SWE is near 80% of median.
- Last month's precipitation for the Powder River Basin was near **75**% of median. Wateryear-to-date precipitation is near **100**% of median.
- The 50% exceedance forecasts for April through July are **below** median (72%) for this basin.






No reservoir data for the basin.








#### **Tongue River Basin**

- The overall Tongue River Basin SWE is near 90% of median.
- Last month's precipitation for the Tongue River Basin was almost 100% of median. Water-year-to-date precipitation is near 115% of median.
- Current reservoir storage is near **100**% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July are **below** median (**81**%) for this basin. Tongue River Reservoir is forecasted to have inflows at **76**% of median.














### Belle Fourche River Basin

- The overall Belle Fourche River Basin SWE is close to **50**% of median.
- Last month's precipitation for the Belle Fourche River Basin was near 50% of median.
  Water-year-to-date precipitation is around 105% of median.
- Current reservoir storage is near **95**% of median for three main reservoirs in the basin.









There are no streamflow forecast points for the basin.





### **Cheyenne River Basin**

- The overall Cheyenne River Basin SWE is near 50% of median.
- Last month's precipitation for the Cheyenne River Basin was between 45 and 50% of median. Water-year-to-date precipitation is around 130% of median.
- $\bullet\,$  Current reservoir storage is near 90% of median for three main reservoirs in the basin.
- The 50% exceedance forecasts for April through July are **below** median (65%) for this basin. Pactola Reservoir inflows are forecasted to be only 60% of median.















# Upper North Platte River Basin

- The overall Upper North Platte River Basin SWE is around 105% of median.
- Last month's precipitation for the Upper North River Basin was near **90**% of median. Water-year-to-date precipitation is around **115**% of median.
- Current reservoir storage is near **80**% of median for two main reservoirs in the basin.
- Streamflow forecasts for April through July are **above** median (**114**%) for this basin. Seminoe Reservoir inflows are forecasted to be **115**% of median.















### Lower North Platte River Basin

- The overall Lower North Platte River Basin SWE is near 80% of median.
- Last month's precipitation for the Lower North Platte River Basin was near 145% of median. Water-year-to-date precipitation is around 120% of median.
- Current reservoir storage is near 100% of median for three main reservoirs in the basin.
- The 50% exceedance forecasts for April through July are **above** median (105%) for this basin. La Prele Creek near Douglas is forecasted to have flows at **85**% of median.















### Laramie River Basin

- The overall Laramie River Basin SWE is around 110% of median.
- Last month's precipitation for the Laramie River Basin was 110 to 115% of median. Water-year-to-date precipitation is near 110% of median.
- Current reservoir storage is around **80**% of median for one main reservoir in the basin.
- Streamflow forecasts for April through July are **above** median (117%) for this basin. Little Laramie River near Filmore is expected to have flows at 117% of median.














#### Sweetwater River Basin

- The overall Sweetwater River Basin SWE is close to 100% of median.
- Last month's precipitation for the Sweetwater River Basin was close to **90**% of median. Water-year-to-date precipitation is near **110**% of median.
- Streamflow forecasts for April through July are **above** median (109%) for this basin.






No reservoir data for the basin.






# South Platte River Basin (WY)

- The overall South Platte River Basin SWE is around **80**% of median.
- Last month's precipitation for the South Platte River Basin was near 170% of median. Water-year-to-date precipitation is close to 100% of median.






No reservoir data for the basin.


There are no streamflow forecast points for the basin.





### Little Snake River Basin

- The overall Little Snake River Basin SWE is near 100% of median.
- Last month's precipitation for the Little Snake River Basin was near 65% of median. Water-year-to-date precipitation is about 110% of median.
- Current reservoir storage is near 50% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July are near the median (95%) for this basin.














### Upper Green River Basin


- The overall Upper Green River Basin SWE is near 100% of median.
- Last month's precipitation for the Upper Green River Basin was between **95** and **100**% of median. Water-year-to-date precipitation is around **115**% of median.
- Current reservoir storage is close to 100% of median for two main reservoirs in the basin.
- Streamflow forecasts for April through July are **below** median (**85**%) for this basin. Green River at Warren Bridge is expected to have flows at **94**% of median.














### Lower Green River Basin

- The overall Lower Green River Basin SWE is near 105% of median.
- Last month's precipitation for the Lower Green River Basin was near 75% of median. Water-year-to-date precipitation is around 120% of median.
- Current reservoir storage is 90 to 95% of median for four main reservoirs in the basin.
- Streamflow forecasts for April through July are **below** median (84%) for this basin.














### Upper Bear River Basin

- The overall Upper Bear River Basin SWE is close to 105% of median.
- Last month's precipitation for the Upper Bear River Basin was near 75% of median. Water-year-to-date precipitation is around 125% of median.
- $\bullet\,$  Current reservoir storage is near 40% of median for two main reservoirs in the basin.
- The 50% exceedance forecasts for April through July are **above** median (**106**%) for this basin. Bear River above Woodruff Narrows Reservoir is expected to have flows at **111**% of median.









# Appendix

#### MEDIAN INFORMATION

#### Transitioning from 1981 – 2010 Averages to 1991 – 2020 Medians

Starting January 2022, the NRCS will use the 30-year **median** as the official normal for snowpack (SWE), precipitation, reservoir storage, and streamflow calculations. The National Water and Climate Center (NWCC) will continue to publish and distribute 30-year <u>averages</u> for alternate normal calculations.

The 30-yr reference period for median and normal calculations has also been recently updated from 1981-2010 to 1991-2020.

Please refer to this NWCC website or more information about the significant changes in data and forecast computations:

https://www.nrcs.usda.gov/wps/portal/wcc/home/snowClimateMonitoring/30YearNormals/

Topics include:

- 1991 2020 Median/Averages Overview
- Calculation Methods
- Differences Between 1991-2020 and Previous Normals
- Median vs. Average
- Retrieving 1991-2020 Normals

For specific seasonal <u>streamflow</u> normal comparisons for NRCS forecasted stations, please refer to:

https://www.wcc.nrcs.usda.gov/ftpref/support/srvo\_norms\_comps/

#### LINKS (for more information/graphics)

#### National Water Climate Center (NWCC)

Interactive maps featuring current conditions of snow, precipitation, reservoir storages:

https://www.nrcs.usda.gov/wps/portal/wcc/home/quicklinks/predefinedMaps/

#### Water Resources Data System and State Climate Office (WRDS)

Clearinghouse of hydrological and climatological data for the State of Wyoming: <u>http://www.wrds.uwyo.edu/</u>

USGS WaterWatch

> Tools and products to monitor streamflow, runoff, drought, and floods:

https://waterwatch.usgs.gov/index.php

# Wyoming Basin Outlook Report National Resources Conservation Service Casper, Wyoming

Issued by:

Released by:

Terry Cosby (Chief) U.S.D.A. Natural Resources Conservation Service Washington D.C. James Bauchert Acting State Conservationist N R C S Casper, Wyoming

# The Following Agencies and Organizations Cooperate with the Natural Resources Conservation Service with Snow Surveys and/or with Data:

#### **FEDERAL:**

United States Department of the Interior (National Park Service)

United States Department of the Interior (Bureau of Reclamation)

United States Department of Agriculture (Forest Service)

United States Department of Commerce NOAA (National Weather Service)

#### STATE:

The Wyoming State Engineer's Office

The University of Wyoming

#### LOCAL:

The City of Cheyenne